




## ANALYTICAL REPORT

### 1-(1,3-diphenylpropan-2-yl)pyrrolidine (C<sub>19</sub>H<sub>23</sub>N)

#### 1-(1,3-diphenylpropan-2-yl)pyrrolidine

Remark – other NPS detected: **none**

Sample ID:	1865-17
Sample description:	crystalline
Sample type:	test purchase /ISF projekt (NFL-SI)
Date of sample receipt (M/D/Y):	10/18/2017
Date of entry (M/D/Y) into NFL database:	11/21/2017
Report updates (if any) will be published here:	<a href="http://www.policija.si/apps/nfl_response_web/seznam.php">http://www.policija.si/apps/nfl_response_web/seznam.php</a>

Substance identified - structure <sup>1</sup> (base form)	
Systematic name	1-(1,3-diphenylpropan-2-yl)pyrrolidine
Other names	
Formula (per base form)	C <sub>19</sub> H <sub>23</sub> N
M <sub>w</sub> (g/mol)	265,4
Salt form/anions detected	HCl
StdInChIKey (per base form)	VQECHRQHFMUVRS-UHFFFAOYSA-N
Other NPS detected	none
Additional info (purity..)	pure by GC-MS, HPLC-TOF

<sup>1</sup> Created by OPSIN free tool: <http://opsin.ch.cam.ac.uk/> DOI: 10.1021/ci100384d

## Report updates

date	comments (explanation)

### Instrumental methods (if applied) in NFL

**1. GC-MS** (Agilent): GC-method is RT locked to tetracosane (9.258 min). Injection volume 1 ml and split mode (1:50). Injector temperature: 280 °C. Chromatographic separation: on column HP1-MS (100% dimethylpolysiloxane), length 30 m, internal diameter 0.25 mm, film thickness 0.25 µm. Carrier gas He: flow-rate 1.2 ml/min. GC oven program: 170 °C for 1 min, followed by heating up to 190 °C at rate 8 °C/min, then heating up to 293 °C at a rate of 18 °C/min, hold for 7.1 min, then heating at 50 °C/min up to 325 °C and finally 6.1 min isothermal. MSD source EI = 70 eV. GC-MS transfer line T= 235 °C, source and quadropole temperatures 280 °C and 180 °C, respectively. Scan range m/z scan range: from 50 (30 until 6 min.) to 550 (300 until 6 min) amu.

**2. HPLC-TOF** (Agilent): 6230B TOF with Agilent 1260 Infinity HPLC with binary pump, column: Zorbax Eclipse XDB-C18, 50 x 4.6 mm, 1.8 micron. Mobile phases (A) 0.1% formic acid and 1mM ammonium formate in water; (B) 0.1% formic acid in methanol (B). Gradient: starting at 5% B, changing to 40% B over 4 min, then to 70% over 2 min and in 5 min to 100%, hold 1 min and back to 5%, equilibration for 1.7 min. The flow rate: 1.0 ml/min; Injection volume 1 µl. MS parameters: 2GHz, Extended Dynamic range mode to a maximum of 1700 amu, acquisition rate 1.30 spectra/sec. Sample ionisation: by Agilent Jet Stream technology (Dual AJS ESI). Ion source: positive ion scan mode with mass scanning from 82 to 1000 amu. Other TOF parameters: drying gas (N<sub>2</sub>) and sheath temperature 325 °C; drying gas flow rate 6 l/min; sheath gas flow rate 8 l/min; nebulizer 25 psig; Vcap. 4000 V; nozzle 2000 V; skimmer 65 V; fragmentor 175 V and Octopole RF 750 V.

**3. FTIR-ATR** (Perkin Elmer): scan range 4000-400 cm<sup>-1</sup>; resolution 4cm<sup>-1</sup>

**4. GC- (MS)-IR** condensed phase (GC-MS (Agilent) & IR (Spectra analyses-Danny)

GC-method: Injection volume 1 ml and split mode (1:5). Injector temperature 280 °C. Chromatographic separation as above (**1**). Split MS : IR = 1: 9.

MSD source EI = 70 eV. GC-MS transfer line T= 235 °C, source and quadropole temperatures 280 °C and 180 °C, respectively. Scan range m/z scan range: from 50 (30 until 6 min.) to 550 (300) amu.

IR (condensed (solid) phase): IR scan range 4000 to 650, resolution 4 cm<sup>-1</sup>.

**5. IC** (anions) (Thermo Scientific, Dionex ICS 2100), Column: IonPac AS19, 2 x 250mm; Eluent: 10mM from 0 to 10 min, 10-58 mM from 10 to 40min; Flow rate: 0.25 ml/min; Temperature: 30 °C; Suppressor: AERS 500 2mm, suppressor current 13mA; Inj. Volume: 25 µl

## Supporting information



Solubility in	result/remark
CH <sub>2</sub> Cl <sub>2</sub>	soluble
MeOH	soluble
H <sub>2</sub> O	soluble

Analytical technique:	applied	remarks
GC-MS (EI ionization)	+	NFL GC-RT (min): 7,74 BP(1): 174; BP(2): 175,BP(3) :91,
HPLC-TOF	+	Exact mass (theoretical): 265,1831; measured value Δppm:-0,46; formula:C19H23N
FTIR-ATR	+	direct measurement (sample as received)
FTIR (condensed phase) always as base form	+	
IC (anions)	+	
NMR (in FKKT)	+	
validation		
other		


# ANALYTICAL RESULTS

MS (EI)


Abundance



## FTIR-ATR - direct measurement (sample as received)



## IR (condensed phase – after chromatographic separation)



# TOF REPORT


<b>Data File</b>	1_1-3-diphenylpropan-2-yl_pyrrolidine-1865-17.d	<b>Sample Name</b>	1865-17
<b>Sample Type</b>	Sample	<b>Position</b>	P2-A4
<b>Instrument Name</b>	6230B TOF LC-MS	<b>User Name</b>	TG
<b>Acq Method</b>	general-19_07_2017-XDB-C18-ESI-final.m	<b>Acquired Time</b>	10/20/2017 9:01:40 AM
<b>IRM Calibration Status</b>	Success	<b>DA Method</b>	Drugs_NFL.m
<b>Comment</b>	MeOH		

## Compound Table


Label	Compound Name	MFG Formula	Obs. RT	Obs. Mass
Cpd 1: 1-(1,3-Diphenylpropan-2-yl)pyrrolidine	1-(1,3-Diphenylpropan-2-yl)pyrrolidine	C19 H23 N	6.62	265.1832

Name	Obs. m/z	Obs. RT	Obs. Mass	DB RT	DB Formula	DB Mass	DB Mass Error (ppm)
1-(1,3-Diphenylpropan-2-yl)pyrrolidine	266.1904	6.62	265.1832	6.62	C19 H23 N	265.1831	-0.46


## Compound Chromatograms



## MFE MS Zoomed Spectrum



## MS Zoomed Spectrum



## MS Spectrum Peak List


Obs. m/z	Charge	Abund	Formula	Ion/Isotope
266.1904	1	27291882	C19 H23 N	(M+H)+
267.1939	1	5925988.01	C19 H23 N	(M+H)+
268.1976	1	509979.21	C19 H23 N	(M+H)+
269.2004	1	32876.92	C19 H23 N	(M+H)+
288.1722	1	2857.18	C19 H23 N	(M+Na)+

--- End Of Report ---

### Peak Integration Report


Sample Name:	1865-17	Inj. Vol.:	25,00
Injection Type:	Unknown	Dilution Factor:	1,0000
Program:	ANIONI	Operator:	kemija
Inj. Date / Time:	20-okt-2017 / 10:36	Run Time:	42,00

No.	Time min	Peak Name	Peak Type	Area $\mu\text{S}\cdot\text{min}$	Height $\mu\text{S}$	Amount mg/L
1,00	9,32	Chloride	BMB	2,27	10,63	n.a.
TOTAL:				2,27	10,63	0,00







## R E P O R T

Contract No.	C1714-17-460078 (Republic of Slovenia, Ministry of the Interior, POLICE)
Sample ID:	<b>1865-17</b>
Received date:	October 26, 2017
Our notebook code:	NFL-1865-17
NMR sample preparation:	18 mg dissolved in 0.7 mL DMSO- <i>d</i> <sub>6</sub>
NMR experiments:	<sup>1</sup> H, <sup>13</sup> C, <sup>1</sup> H- <sup>1</sup> H <i>gs</i> -COSY, <sup>1</sup> H- <sup>13</sup> C <i>gs</i> -HSQC, <sup>1</sup> H- <sup>13</sup> C <i>gs</i> -HMBC, <sup>1</sup> H- <sup>15</sup> N <i>gs</i> -HMBC
Proposed structure, formula, exact mass, molecular weight:	 <p>Chemical Formula: C<sub>19</sub>H<sub>24</sub>N<sup>+</sup> Exact Mass: 266,1903 Molecular Weight: 266,4075</p>
Chemical name:	1-(1,3-Diphenylpropan-2-yl)pyrrolidin-1-ium ion
Comments:	- Structure elucidation based on 1D and 2D NMR spectra and HRMS.
Supporting information:	Copies of <sup>1</sup> H and <sup>13</sup> C NMR spectra
Authors:	Martin Gazvoda, Marko Krivec, Janez Košmrlj
Date of report:	November 13, 2017







```

Current
NAME
EXPNO
PROCNO
F2 - Acq
Date_
Time_
INSTRUM
PROBHD
PULPROG
TD
SOLVENT
NS
DS
SWH
FIDRES
AQ
RG
DW
DE
TE
D1
D11
TD0
=====
SFO1
NUC1
P1
PLM1
=====
SFO2
NUC2
CEPRG1
PCPD2
PLM2
PLM12
PLM13
F2 - Prc
SI
SF
WDW
SSB
TB
GB
PC

```