


ANALYTICAL REPORT

Norflurazepam (Norfludiazepam) (C₁₅H₁₀ClFN₂O)

7-chloro-5-(2-fluorophenyl)-2,3-dihydro-1H-1,4-benzodiazepin-2-one

Remark – other NPS detected: none

Sample ID:	2188-20
Sample description:	powder
Sample type:	test purchase /ISF projekt (NFL-SI)
Date of entry (DD/MM/YYYY) into NFL database:	20/10/2020
Report updates (if any) will be published here:	http://www.policija.si/apps/nfl_response_web/seznam.php

Substance identified - structure ¹ (base form)	
Systematic name	7-chloro-5-(2-fluorophenyl)-2,3-dihydro-1H-1,4-benzodiazepin-2-one
Other names	Norfludiazepam; N-Desalkylflurazepam; Descarbethoxyloflazepate; N-Desalkyl-2-oxoquazepam;
Formula (per base form)	C ₁₅ H ₁₀ ClFN ₂ O
M _w (g/mol)	288,71
Salt form/anions detected	base
StdInChIKey (per base form)	UVCOILFBWYKHHB-UHFFFAOYSA-N
Other NPS detected	none
Additional info (purity..)	>95% purity of a sample based on 1H NMR

¹ Created by OPSIN free tool: <http://opsin.ch.cam.ac.uk/> DOI: 10.1021/ci100384d

Report updates

date	comments (explanation)

Instrumental methods (if applied) in NFL

1. GC-MS (Agilent): GC-method is RT locked to tetracosane (9.258 min). Injection volume 1 µl and split mode (1:50). Injector temperature: 280 °C. Chromatographic separation: on column HP1-MS (100% dimethylpolysiloxane), length 30 m, internal diameter 0.25 mm, film thickness 0.25 µm. Carrier gas He: flow-rate 1.2 ml/min. GC oven program: 170 °C for 1 min, followed by heating up to 190 °C at rate 8 °C/min, then heating up to 293 °C at a rate of 18 °C/min, hold for 7.1 min, then heating at 50 °C/min up to 325 °C and finally 6.1 min isothermal. MSD source EI = 70 eV. GC-MS transfer line T= 235°C, source and quadropole temperatures 280°C and 180°C, respectively. Scan range m/z scan range: from 50 (30 until 6 min.) to 550 (300 until 6 min) amu.

2. HPLC-TOF (Agilent): 6230B TOF with Agilent 1260 Infinity HPLC with binary pump, column: Zorbax Eclipse XDB-C18, 50 x 4.6 mm, 1.8 micron. Mobile phases (A) 0.1% formic acid and 1mM ammonium formate in water; (B) 0.1% formic acid in methanol (B). Gradient: starting at 5% B, changing to 40% B over 4 min, then to 70% over 2 min and in 5 min to 100%, hold 1 min and back to 5%, equilibration for 1.7 min. The flow rate: 1.0 ml/min; Injection volume 1 µl. MS parameters: 2GHz, Extended Dynamic range mode to a maximum of 1700 amu, acquisition rate 1.30 spectra/sec. Sample ionisation: by Agilent Jet Stream technology (Dual AJS ESI). Ion source: positive ion scan mode with mass scanning from 82 to 1000 amu. Other TOF parameters: drying gas (N2) and sheath temperature 325 °C; drying gas flow rate 6 l/min; sheath gas flow rate 8 l/min; nebulizer 25 psig; Vcap. 4000 V; nozzle 2000 V; skimmer 65 V; fragmentor 175 V and Octopole RF 750 V.

3. FTIR-ATR (Perkin Elmer): scan range 4000-400 cm⁻¹; resolution 4cm⁻¹

4. GC- (MS)-IR condensed phase (GC-MS (Agilent) & IR (Spectra analyses-Danny)

GC-method: Injection volume 1 µl and split mode (1:5). Injector temperature 280 °C. Chromatographic separation as above (1). Split MS : IR = 1: 9.

MSD source EI = 70 eV. GC-MS transfer line T= 235°C, source and quadropole temperatures 280°C and 180°C, respectively. Scan range m/z scan range: from 50 (30 until 6 min.) to 550 (300) amu.

IR (condensed (solid) phase): IR scan range 4000 to 650, resolution 4 cm⁻¹.

5. IC (anions) (Thermo Scientific, Dionex ICS 2100), Column: IonPac AS19, 2 x 250mm; Eluent: 10mM KOH from 0 to 10 min, 10-58 mM from 10 to 40min; Flow rate: 0.25 ml/min; Temperature: 30°C; Suppressor: AERS 500 2mm, suppressor current 13mA; Inj. Volume: 25 µl

Supporting information


Solubility in	result/remark
CH ₂ Cl ₂	soluble
MeOH	soluble
H ₂ O	not soluble

Analytical technique:	applied	remarks
GC-MS (EI ionization)	+	NFL GC-RT (min): 9,42 BP(1): 259; BP(2): 260,BP(3) :287,
HPLC-TOF	+	Exact mass (theoretical): 288,0466; measured value Δppm:-1,7; formula:C15H10ClFN2O
FTIR-ATR	+	direct measurement (sample as received)
FTIR (solid phase) always as base form	+	
IC (anions)	+	
NMR (in FKKT)		
validation		
other		


ANALYTICAL RESULTS

MS (EI)


Abundance



FTIR-ATR - direct measurement (sample as received)



IR (solid phase – after chromatographic separation)



TOF REPORT


Data File	Norflurazepam_2188-20.d	Sample Name	ID-2188-20
Sample Type	Sample	Position	P2-A5
Instrument Name	6230B TOF LC-MS	User Name	
Acq Method	general-15_01_2020-XDB-C18-ESI+.m	Acquired Time	9/2/2020 8:30:40 PM
IRM Calibration Status	Success	DA Method	a-Drugs_NFL.m
Comment			

Compound Table


Label	Compound Name	MFG Formula	Obs. RT	Obs. Mass
Cpd 1: Norflurazepam	Norflurazepam	C15 H10 Cl F N2 O	7.6	288.0471

Name	Obs. m/z	Obs. RT	Obs. Mass	DB RT	DB Formula	DB Mass	DB Mass Error (ppm)
Norflurazepam	289.0543	7.6	288.0471	7.6	C15 H10 Cl F N2 O	288.0466	-1.7


Compound Chromatograms



MFE MS Zoomed Spectrum



MS Zoomed Spectrum



MS Spectrum Peak List


Obs. m/z	Charge	Abund	Formula	Ion/Isotope
289.0543	1	21653206	C15 H10 Cl F N2 O	(M+H)+
290.0574	1	3767868.01	C15 H10 Cl F N2 O	(M+H)+
291.0518	1	7011522.31	C15 H10 Cl F N2 O	(M+H)+
292.0547	1	1146125.02	C15 H10 Cl F N2 O	(M+H)+
311.0364	1	839162.25	C15 H10 Cl F N2 O	(M+Na)+
313.0335	1	262982.51	C15 H10 Cl F N2 O	(M+Na)+
599.0828	1	1603839	C15 H10 Cl F N2 O	(2M+Na)+
600.0861	1	554269.09	C15 H10 Cl F N2 O	(2M+Na)+
601.081	1	1110067.06	C15 H10 Cl F N2 O	(2M+Na)+
602.0832	1	350243.27	C15 H10 Cl F N2 O	(2M+Na)+

--- End Of Report ---

Peak Integration Report

Sample Name:	2188_IC	Inj. Vol.:	25,00
Injection Type:	Unknown	Dilution Factor:	1,0000
Program:	ANIONI	Operator:	kemija
Inj. Date / Time:	02-sep-2020 / 13:52	Run Time:	43,00

No.	Time min	Peak Name	Peak Type	Area $\mu\text{S}\cdot\text{min}$	Height μS	Amount n.a.
TOTAL:				0,00	0,00	0,0




University
of Ljubljana


Faculty of Chemistry
and Chemical Technology



R E P O R T

Contract No.	C1714-19-460155 (Republic of Slovenia, Ministry of the Interior, POLICE)
Sample ID:	2188-20
Received date:	September 10, 2020
Our notebook code:	NFL-2188-20
NMR sample preparation:	19.7 mg dissolved in 0.7 mL DMSO- <i>d</i> ₆
NMR experiments:	¹ H, ¹³ C, ¹ H- ¹ H <i>gs</i> -COSY, ¹ H- ¹³ C <i>gs</i> -HSQC, ¹ H- ¹³ C <i>gs</i> -HMBC, ¹ H- ¹⁵ N <i>gs</i> -HMBC, ¹⁹ F
Proposed structure with formula, exact mass, molecular weight:	 <p>Chemical Formula: C₁₅H₁₀ClFN₂O Exact Mass: 288.04657 Molecular Weight: 288.70640</p>
Chemical name:	7-chloro-5-(2-fluorophenyl)-1,3-dihydro-2H-benzo[e][1,4]diazepin-2-one
Comments:	- Structure elucidation based on 1D and 2D NMR spectra and HRMS. ->95% purity of a sample based on ¹ H NMR spectrum.
Supporting information:	Copies of ¹ H and ¹³ C NMR spectra, ¹ H and ¹³ C FIDs.
Principal investigator:	Prof. Dr. Janez Košmrlj
Date of report:	October 15, 2020

NFL-2188-20
1H



Current Data Parameters
NAME NFL-2188-20
EXPNO 1
PROCNO 1


F2 - Acquisition Parameters

Date_ 20201008
Time_ 1.19
INSTRUM 5 mm PABBO BB-
PROBHD spect
PULPROG zg30
TD 65536
SOLVENT DMSO
NS 16
DS 2
SWH 10000.000 Hz
FIDRES 0.152588 Hz
AQ 3.276799 sec
RG 101
DW 50.000 usec
DE 6.50 usec
TE 296.0 K
D1 1.0000000 sec
TD0 1

==== CHANNEL f1 =====
SFO1 500.130885 MHz
NUC1 1H
P1 8.70 usec
PLW1 26.0000000 W

F2 - Processing parameters
SI 65536
SF 500.1300044 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

NFL-2188-20
13C



Current Data Parameters
NAME NFL-2188-20
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20201008
Time 4.02

INSTRUM spect
PROBHD 5 mm PABBO BB-
PULPROG zgpg30
TD 65536
SOLVENT DMSO
NS 3072
DS 4
SWH 29761.904 Hz
FIDRES 0.454131 Hz
AQ 1.1010048 sec
RG 2050
DM 16.800 usec
DE 6.50 usec
TE 296.0 K
D1 2.0000000 sec
D11 0.03000000 sec
TD0 1

==== CHANNEL f1 =====

SFO1 125.7703637 MHz
NUC1 13C
P1 8.70 usec
PLW1 122.0000000 W

==== CHANNEL f2 =====

SFO2 500.1320005 MHz
NUC2 1H
CPDPRG2 waltz16
PCPD2 80.00 usec
PLW2 26.00000000 W
PLW12 0.30046001 W
PLW13 0.15113001 W

F2 - Processing parameters

SI 32768
SF 125.7578442 MHz
WDW EM
SSB 0
IB 1.00 Hz
GB 0
PC 1.40